

Negative heat capacity and non-extensive kinetic theory

R. Silva*

*Departamento de Física, Universidade do Estado do Rio
Grande do Norte, 59610-210, Mossoró - RN, Brazil*

J. S. Alcaniz†

*Astronomy Department, University of Washington,
Seattle, Washington, 98195-1580, USA and
Departamento de Física, Universidade Federal do Rio Grande do Norte,
C.P. 1641, Natal - RN, 59072-970, Brazil*

(Dated: February 2, 2008)

Abstract

The negative nature of the heat capacity C_V of thermodynamically isolated self-gravitating systems is rediscussed in the framework of a non-extensive kinetic theory. It is found that the dependence of C_V on the non-extensive parameter q gives rise to a negative branch with the critical value corresponding to $q = 5/3$ ($C_V \rightarrow -\infty$).

PACS numbers:

*Electronic address: rsilva@uern.br

†Electronic address: alcaniz@dfte.ufrn.br

Consider a simple analogy between ideal gas and self-gravitating system. The temperature of this system is defined by

$$\frac{1}{2}mv^2 = \frac{3}{2}k_B T, \quad (1)$$

where m is the mass of a particle (e.g., a star) and k_B is the Boltzmann's constant. If such a system is composed by N particles, its total kinetic energy is $K = \frac{3}{2}Nk_B T$ which, according to the virial theorem, is equal to minus its total energy, i.e., $E = -K$. Therefore, the heat capacity of the system is

$$C_V = \frac{dE}{dT} = -\frac{3}{2}Nk_B, \quad (2)$$

which is clearly a negative quantity. In other words, this means that by losing energy the system automatically grows hotter, or still, that a self-gravitating system can never be in a true thermodynamical equilibrium state (see [1] and references therein). In actual fact, it is well known that when discussed in the context of the microcanonical ensemble a self-gravitating system interacting via long range forces exhibits a negative heat capacity [2]. The negativeness of the heat capacity has been largely studied in many astrophysical phenomena. For example, Lynden-Bell & Wood [3] used such a concept to explain the gravothermal catastrophe for globular clusters while Beckenstein [4] and Hawking [5] used it in the context of black hole thermodynamics.

On the other hand, a considerable theoretical effort has been concentrated on the study of the statistical description of a large variety of physical systems, resulting in the extension of the Boltzmann-Gibbs' statistical mechanics. Particular examples are systems endowed with long duration memory, anomalous diffusion, turbulence in pure-electron plasma, self-gravitating systems or, more generally, systems endowed with long range interactions. In order to deal with such problems, Tsallis [6] proposed the following generalization of the Boltzmann-Gibbs (BG) entropy formula

$$S_q = -k_B \frac{1 - \sum_i p_i^q}{(q-1)}, \quad (3)$$

where p_i is the probability of the i^{th} microstate and q is a parameter quantifying the degree of non-extensivity. In the limit $q \rightarrow 1$ the celebrated BG extensive formula, $S_1 = -k_B \sum_i p_i \ln p_i$, is readily recovered as a particular case.

In the astrophysical context, the first applications of this non-extensive q -statistics studied stellar polytropes [7] and the peculiar velocity function of galaxy clusters [8]. The Jeans

gravitational instability criterion for a collisionless system was also recently rediscussed in the context of this enlarged formalism [9]. In this latter study, it was shown that such systems present instability even for wavelengths of the disturbance smaller than the standard critical Jeans value. Recently, Taruya & Sakagami [10, 11] discussed a stability problem in the microcanonical ensemble by extending Padmanabhan's [2] classical analysis of the Antonov instability to the case of polytropic distributions functions. More recently, Chavanis [12, 13, 14] considered the Tsallis' entropy as a particular H -function corresponding to isothermal stellar systems and stellar polytropes. In this approach, the maximization of the H -function at fixed mass and energy reveals naturally a thermodynamical analogy with the study of the dynamical stability in collisionless stellar systems [24].

In this *Letter*, we discuss the negative nature of the specific capacity C_V of self-gravitating systems in the framework of a kinetic theory resulting from this non-extensive formalism. To do so, we consider self-gravitating systems that can be treated as isolated systems in a first approximation, e.g., globular cluster [15] or elliptical galaxies [16]. We derive a new analytical expression for this quantity which gives rise to an entire negative branch. We show that for this kind of system the value $q = 5/3$ is an upper limit for the non-extensive parameter. In connection with stellar polytropes, the value of the non-extensive parameter $q = 5/3$ corresponding to a polytropic index $n = 3$ or $n = -1$, depending on the expression between n and q .

It has been shown that the kinetic foundations of the Tsallis' thermostatistic lead to a velocity distribution for free particles given by [17, 18]

$$f_0(v) = B_q \left[1 - (1 - q) \frac{mv^2}{2k_B T} \right]^{1/(1-q)}. \quad (4)$$

In this velocity q -distribution, the q -parameter quantifies the nonadditivity property of the associated gas entropy whose the main effect at the level of the distribution is to replace the standard Gaussian form by a power law. Mathematically, this result follows directly from the well known identity $\lim_{w \rightarrow 0} (1 + wx)^{1/w} = \exp(x)$ [19]. The quantity B_q is a q -dependent normalization constant whose expression for $q \leq 1$ and $q \geq 1$ are respectively given by

$$\frac{B_{q \leq 1}}{B_{q \geq 1}} = \begin{cases} n(1 - q)^{1/2} \left(\frac{5-3q}{2} \right) \left(\frac{3-q}{2} \right) \frac{\Gamma(\frac{1}{2} + \frac{1}{1-q})}{\Gamma(\frac{1}{1-q})} \left(\frac{m}{2\pi k_B T} \right)^{3/2} \\ n(q - 1)^{3/2} \frac{\Gamma(\frac{1}{q-1})}{\Gamma(\frac{1}{q-1} - \frac{3}{2})} \left(\frac{m}{2\pi k_B T} \right)^{3/2} \end{cases}, \quad (5)$$

where T is the temperature and n is the particle number density. This non-extensive velocity

distribution can be derived at least from two different methods: (i) through a simple non-extensive generalization of the Maxwell ansatz, $f(v) \neq f(v_x)f(v_y)f(v_z)$, which follows from the introduction of statistical correlations between the components of the velocities [17] and (ii) by using a more rigorous treatment based on the non-extensive formulation of the Boltzmann H -theorem, which requires $q > 0$ [18].

In order to investigate the q -dependence of the specific capacity, we first consider a cloud of ideal gas within the nonrelativistic gravitational context, which is analog to a self-gravitating collisionless gas. The kinetic energy of this system is simply given by $K = \frac{1}{2}m \langle v^2 \rangle$, with the statistical content included in the average square velocity of the particles $\langle v^2 \rangle$. In this way, the non-extensivity is introduced through a new derivation of expectation value [20]

$$\langle v^2 \rangle_q = \frac{\int_{-v_m}^{v_m} f^q v^2 d^3v}{\int_{-v_m}^{v_m} f^q d^3v}, \quad (6)$$

where $v_m = \left(\frac{2k_B T}{m(1-q)} \right)^{1/2}$ is a thermal cutoff on the maximum value allowed for the velocity of the particles ($q < 1$), whereas for the power law q -distribution without cutoff ($q > 1$) $v_m \rightarrow \infty$. This q -expectation value can be easily evaluated for $q \neq 1$, and the result corresponds to a q -parameterized family of the square velocity of the particles $\langle v^2 \rangle_q$ given by

$$\langle v^2 \rangle_q = \frac{6}{5-3q} \frac{k_B T}{m}, \quad \text{for } q < 5/3, \quad (7)$$

with $\langle v^2 \rangle = \frac{3k_B T}{m}$ being the classical value easily obtained in the limit $q \rightarrow 1$ (It is worth mentioning that the virial theorem is not modified in this non-extensive formalism [21]).

Now, by substituting the non-extensive q -expectation value given by Eq. (7) into the heat capacity relation Eq. (2) one finds

$$C_V = -\frac{3}{5-3q} N k_B, \quad (8)$$

which clearly reduces to the Maxwellian limit $C_V = -\frac{3}{2} N k_B$ for $q = 1$. To better understand the physical implications of the above expression, in Fig. 1 we plot the quantity C_V/Nk_B as a function of the non-extensive parameter q . The graph clearly shows that the specific capacity is a negative quantity for the interval $q < 5/3$. For $q = 5/3$ (critical value) the specific capacity diverges ($C_V \rightarrow -\infty$). From this analysis, it is possible to conclude that this particular q -dependence of C_V leads to the classical gravothermal catastrophe ($C_V < 0$) only for power laws with non-extensive parameter in the range $0 < q < 5/3$.




FIG. 1: The quantity C_V/Nk_B for a self-gravitating collisionless gas obeying the non-extensive Tsallis' q -statistic is shown as a function of the non-extensive parameter q . We see that there is a negative branch for the interval $0 < q < 5/3$. For $q = 5/3$ the specific capacity diverges ($C_V \rightarrow -\infty$).

At this point we emphasize that the true nature of the non-extensive q -parameter appearing in the Tsallis' statistical framework remains as a completely open question at present. In this way, it is important to show empirical manifestations of these non-extensive effects. In this *Letter* we have discussed a clear signature of non-extensive effects on some kind of self-gravitating systems. In particular, we have shown that the q -dependence of the specific capacity for these systems is physically consistent only for power-laws lying in the interval $0 < q < 5/3$. This particular range of values for the non-extensive parameter q is less restrictive than that one obtained by Abe [22]. We suspect that such a difference occurs because the Abe's approach is addressed in the context of the canonical ensemble and, as has been shown elsewhere, in this ensemble self-gravitating systems do not present negative specific heat [2]. We also note that, when identified with the polytropic index $n = \frac{3}{2} + \frac{1}{q-1}$ [7], the critical value $q = 5/3$ is identical to the one that gives rises to the gravitational stability/instability transition, i.e., $n = 3$ [12] (or $n = -1$ if $n = \frac{1}{2} + \frac{1}{1-q}$ [11]). In a couple of papers by Taruya & Sakagami [11, 23] it was shown that a thermodynamic instability

appears for values of $n > 5$ in systems confined in adiabatic walls (microcanonical ensemble) and for values of $n > 3$ in systems surrounded by a thermal bath. In fact, a more consistent study of the gravothermal catastrophe must consider a self-gravitating ideal gas contained in a spherical container. We shall return to this discussion in a more complete analysis involving general non-extensive kinetic theories which will appear in a forthcoming communication.

Acknowledgments: The authors are very grateful to J. A. S. Lima, P. H. Chavanis and A. Taruya for helpful discussions and a critical reading of the manuscript. This work is supported by the project CNPq (62.0053/01-1-PADCT III/Milenio). JSA is also supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq - Brasil).

-
- [1] Lynden-Bell, *Physica A* 263 (1998) 293
 - [2] T. Padmanabhan, *Phys. Rep.* 188 (1990) 285.
 - [3] D. Lynden-Bell, R. Wood, *Mon. Not. R. Soc.* 136 (1967) 101.
 - [4] J. D. Beckenstein, *Phys. Rev. D* 9 (1974) 3292.
 - [5] S. W. Hawking, *Nature* 248 (1974) 30.
 - [6] C. Tsallis, *J. Stat. Phys.* 52 (1988) 479.
 - [7] A. Plastino, A. R. Plastino, *Phys. Lett. A* 174 (1993) 384.
 - [8] A. Lavagno, G. Kaniadakis, M. Rego-Monteiro, P. Quarati, C. Tsallis, *Astroph. Lett. and Comm.* 35 (1998) 449.
 - [9] J. A. S. Lima, R. Silva, J. Santos, *A&A* 396 (2002) 309.
 - [10] A. Taruya, M. Sakagami, *Physica A* 318 (2003) 387.
 - [11] A. Taruya, M. Sakagami, *Physica A* 322 (2003) 285.
 - [12] P. H. Chavanis, *A&A* 386 (2002) 732.
 - [13] P. H. Chavanis, *A&A* 401 (2003) 15.
 - [14] P. H. Chavanis, C. Sire, *cond-mat/0303088*.
 - [15] D. Lynden-Bell, R. Wood, *Mon. Not. Roy. Astron. Soc.* 138 (1968) 495.
 - [16] J. Hjorth, J. Madsen, *Mon. Not. Roy. Astron. Soc.* 265 (1993) 237.
 - [17] R. Silva, A. R. Plastino, J. A. S. Lima, *Phys. Lett. A* 249 (1998) 401.
 - [18] J. A. S. Lima, R. Silva, A. R. Plastino, *Phys. Rev. Lett.* 86 (2001) 2938.
 - [19] M. Abramowitz, I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publ., New York, 1972) p. 257.
 - [20] C. Tsallis, R. S. Mendes, A. R. Plastino, *Physica A* 261 (1998) 534.
 - [21] S. Martínez, F. Pennini, A. Plastino, *cond-mat/0006139*.
 - [22] S. Abe, *Phys. Lett. A* 263 (1999) 424.
 - [23] A. Taruya, M. Sakagami, *Phys. Rev. Lett.* 90 (2003) 181101.
 - [24] For a complete and updated list of references see <http://tsallis.cat.cbpf.br/biblio.htm>